## NOVELLO FREQUENCY CONTROL FOR A 220 GHZ RADIOMETER WITH A 110 GHZ GUNN OSCILLATOR

Olga Borie-Lubecke\*, Richard F. Denning, Michael A. Janssen and Margaret A. 1 rerking

Jet Propulsion 1 aboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 911 09-8099, USA

Abstract - Because of the drive for "faster, belter, cheaper" missions, future space missions will require low-power, compact millimeter and submillimeter wave instruments. Eliminating phase-locked loop (PLL) systems as well as active thermal control systems from standard receivers can result in a significant reduction of mass and power in some applications, such as planetary atmospheric sounding. 1 lere we report on a novel 1.0 frequency control for millimeter and submillimeter wave subharmonic Schottky receivers using GUNN oscillators for the 1.(). In (his approach, the GUNN oscillator frequency is voltage tuned to compensate for the frequency drift due to the temperature change, and therefore neither a 1)1.1, nor active thermal control are required. A temperature-compensated, fixed-luncd GUNN oscillator was characterized for a wide temperature range (0 to 50 °C), and frequency versus voltage and temperature behavior was recorded. Based on this information, a simple computer program was written to calculate the required bias voltage for a desired frequency and a given temperature. A 220 G] lz subharmonic receiver using such an 1.0 frequency control was tested with a laboratory signal source. It was shown that the GUNN frequency could be repeatably predicted within a few MHz, giving sufficient accuracy for Mars atmospheric sounding.

W c would like to thank R. J. Dengler, '1'. C. Koch, M. J. Mahoney, D. O. Muhleman, J. E. Oswald, 1'.11. Siegel, and U. Zimmerman for valuable (discussions, and A. B. Hubbard for technical support. This research was sponsored by the National Aeronauties and Space Administration, Office of Space Sciences.

\*Olga Boric-Lubecke was an NRC Resident Research Associate at JPL. She is currently with The Institute of Physical and Chemical Research, Sendai, Japan.